



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 45/06, 31/485	A1	(11) International Publication Number: WO 99/45963 (43) International Publication Date: 16 September 1999 (16.09.99)
(21) International Application Number: PCT/US99/05336 (22) International Filing Date: 11 March 1999 (11.03.99) (30) Priority Data: 09/042,248 13 March 1998 (13.03.98) US (71) Applicant: ALGOS PHARMACEUTICAL CORPORATION [US/US]; Collingwood Plaza, 4900 Route 33, Neptune, NJ 07753 (US). (72) Inventor: CARUSO, Frank, S.; 2 Bowling Green, Colts Neck, NJ 07722 (US). (74) Agents: DILWORTH, Peter, G. et al.; Dilworth & Barrese, 333 Earle Ovington Boulevard, Uniondale, NY 11553 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: ANALGESIC COMBINATION COMPRISING NMDA RECEPTOR ANTAGONISTS AND NARCOTIC ANALGESICS (57) Abstract <p>The analgesic effectiveness of a narcotic agonist-antagonist analgesic is significantly potentiated by administering a narcotic agonist-antagonist analgesic together with at least one nontoxic NMDA receptor antagonist. The narcotic agonist-antagonist analgesic is selected from the group consisting of pentazocine, nalbuphine, butorphanol, buprenorphine, meptazinol, dezocine, nalorphine, cyclazocine and pharmaceutically acceptable salts thereof. The nontoxic NMDA receptor antagonist is at least one member selected from the group consisting of dextromethorphan, dextrophan, amantadine, memantine and pharmaceutically acceptable salts thereof.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ANALGESIC COMBINATION COMPRISING NMDA RECEPTOR ANTAGONISTS AND NARCOTIC ANALGESICS

BACKGROUND OF THE INVENTION

5 This invention relates to an analgesic composition and method for alleviating pain. More particularly, this invention is concerned with alleviating pain by administering to a mammal exhibiting pain (a) an analgesia-effective amount of at least one narcotic agonist-antagonist analgesic; and, (b) a narcotic agonist-antagonist analgesic-potentiating amount of at least one nontoxic antagonist, or
10 blocker, for the N-methyl-D-aspartate (NMDA) receptor.

 Narcotic agonist-antagonist analgesics are known. In general, narcotic agonist-antagonist analgesics constitute a distinct subclass of opioids and are differentiated from the latter by their mixed actions, meaning, they are not full agonists at all opioid receptors, e.g., μ , δ , κ , etc. receptors. Instead, narcotic
15 agonist-antagonist analgesics are believed to either exert their analgesic action by working as agonists at some opioid receptors and antagonists or very weak agonists at other opioid receptors, i.e., mixed narcotic agonist-antagonists, or exert their analgesic action by working as agonists at some opioid receptors, i.e., partial agonists. Mixed narcotic agonist-antagonist analgesics will typically be the
20 combination of μ antagonism coupled with κ agonism. Partial agonist analgesics will typically be μ agonism.

 Heretofore, there has been no recognition or appreciation that the analgesic effectiveness of a narcotic agonist-antagonist analgesic can be appreciably potentiated, or enhanced, by administration of a narcotic agonist-antagonist
25 analgesic prior to, with or following the administration of a narcotic agonist-antagonist analgesic-potentiating amount of a nontoxic NMDA receptor antagonist.

SUMMARY OF THE INVENTION

In accordance with the present invention, an analgesic composition for alleviating pain is provided which comprises (a) an analgesia-effective amount of at least one narcotic agonist-antagonist analgesic; and, (b) a narcotic agonist-antagonist analgesic-potentiating amount of at least one nontoxic N-methyl-D-aspartate receptor antagonist.

Further in accordance with the present invention, a method of alleviating pain is provided which comprises administering to a mammal exhibiting pain (a) an analgesia-effective amount of at least one narcotic agonist-antagonist analgesic; and, (b) a narcotic agonist-antagonist analgesic-potentiating amount of at least one nontoxic N-methyl-D-aspartate receptor antagonist.

The analgesic composition and method of alleviating pain of this invention are applicable to the treatment of all varieties of pain, e.g., arthritic pain, lumbosacral pain, musculo-skeletal pain, pain associated with a sore throat, etc. Thus, for a mammal in need of pain relief, an enhanced level of analgesia for an equivalent dosage of at least one narcotic agonist-antagonist analgesic, or an equivalent level of analgesia for a reduced dosage of at least one narcotic agonist-antagonist analgesic, can be achieved when at least one narcotic agonist-antagonist analgesic is administered prior to, with or following the administration of the nontoxic NMDA receptor antagonist.

The term "narcotic agonist-antagonist analgesic" shall be understood as referring to those analgesics which exhibit mixed narcotic agonist-antagonist or partial agonist properties.

The term "N-methyl-D-aspartate receptor" shall be understood to include all of the binding site subcategories associated with the NMDA receptor, e.g., the glycine-binding site, the phenylcyclidine (PCP)-binding site, etc., as well as the NMDA channel. Thus, the invention herein contemplates the use of nontoxic

substances that block an NMDA receptor binding site, e.g., dextromethorphan, or block the NMDA channel, e.g., a source of magnesium such as magnesium sulfate.

The term "nontoxic" as used herein shall be understood in a relative sense and is intended to designate any substance that has been approved by the
5 United States Food and Drug Administration ("FDA") for administration to humans or, in keeping with established regulatory criteria and practice, is susceptible to approval by the FDA for administration to humans. The term "nontoxic" is also used herein to distinguish the NMDA receptor antagonists, or blockers, that are useful in the practice of the present invention from NMDA receptor antagonists
10 such as MK 801 (the compound 5-methyl-10,11-dihydro-SH-dibenze[a,d] cyclohepten-5,10-imine), CPP (the compound 3-[2-carboxypiperazin-4-yl] propyl-1-phosphonic acid) and PCP (the compound 1-(1-phenylcyclohexyl)piperidine) whose toxicities effectively preclude their therapeutic use.

The terms "potentiate" and "potentiating" are used herein in their art-
15 recognized sense, i.e., as referring to a significant increase in the level of pain-alleviating activity for the combination of narcotic agonist-antagonist analgesic and nontoxic NMDA receptor antagonist compared with that which could have been expected based on the activities of the narcotic agonist-antagonist analgesic administered alone and nontoxic NMDA receptor antagonist administered alone.

20 The term "pain-alleviating" shall be understood herein to include the expressions "pain-suppressing" and "pain-inhibiting" as the invention is applicable to the alleviation of existing pain as well as the suppression or inhibition of pain which would otherwise ensue from an imminent pain-causing event.

The expression "analgesia-effective amount" as applied to the
25 narcotic agonist-antagonist analgesic employed in the composition and method of this invention shall be understood to mean an amount of narcotic agonist-antagonist analgesic which when administered by itself or in combination with the nontoxic NMDA receptor antagonist provides significant analgesic activity.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

Any of the narcotic agonist-antagonist analgesics heretofore used to alleviate pain can be used herein. For listings of narcotic agonist-antagonist analgesics, see, e.g., Goodman and Gilman's "The Pharmaceutical Basis of Therapeutics", 8th ed., McGraw-Hill, Inc. (1990), pp. 510-514 and "Remington's Pharmaceutical Sciences", 17th ed., Mack Publishing Company (1985), pp. 1099-1110. Specific narcotic agonist-antagonist analgesics that can be used herein include pentazocine, pentazocine hydrochloride, nalbuphine, nalbuphine hydrochloride, butorphanol, butorphanol tartrate, buprenorphine, buprenorphine hydrochloride, meptazinol, dezocine, nalorphine, cyclazocine, their pharmaceutically acceptable salts and the like.

Among the nontoxic substances that block the NMDA receptor and as such are useful for potentiating the analgesic activity of the foregoing narcotic agonist-antagonist analgesics in accordance with this invention are dextromethorphan ((+)-3-hydroxy-N-methylmorphinan), its metabolite dextrorphan ((+)-3-hydroxy-N-methylmorphinan), amantadine (1-amino adamantine), memantine (3,5 dimethylaminoadamantone), their mixtures and their pharmaceutically acceptable salts. Other useful nontoxic substances that block the NMDA receptor include pyrroloquinoline quinone, 4-hydroxy-2(1H)-quinolone derivatives and cis-4-(phosphono-methyl)-2-piperidinecarboxylic acid. Of the foregoing nontoxic substances that block the NMDA receptor, dextromethorphan in the form of its hydrobromide salt is preferred for use herein due to its ready availability and its established use in over-the-counter medications where it functions as a cough suppressant.

With regard to dosage levels, the narcotic agonist-antagonist analgesic must be present in an analgesia-effective amount, e.g., at a level corresponding to the generally recommended adult human dosages for a particular narcotic agonist-antagonist analgesic, and the nontoxic NMDA receptor antagonist

must be present at a level that potentiates the effectiveness of the narcotic agonist-antagonist analgesic. Specific dosage levels for the narcotic agonist-antagonist analgesic that can be used herein as given, inter alia, in the "Physicians' Desk Reference", 1997 Edition (Medical Economics Data Production Company, Montvale, NJ) as well as in other reference works including Goodman and Gilman's "The Pharmaceutical Basis of Therapeutics" and "Remington's Pharmaceutical Sciences" both of which as referred to above. Given the wide variation in dosage level of the narcotic agonist-antagonist analgesic which depends to a large extent on the specific narcotic agonist-antagonist analgesic being administered, there can similarly be a wide variation in the dosage level of the nontoxic NMDA receptor antagonist. These amounts can be determined for a particular drug combination in accordance with this invention employing routine experimental testing. In the case of pentazocine and the nontoxic NMDA receptor antagonist dextromethorphan, dosages of from about 30 to about 250 mg/day of the former coadministered with from about 30 to about 120 mg/day of the latter will usually provide acceptable results.

While the narcotic agonist-antagonist analgesic and potentiating nontoxic NMDA receptor antagonist need not be administered together, they must both be present in the patient at effective levels at the same time. While it is within the scope of the invention to separately administer the narcotic agonist-antagonist analgesic and the nontoxic NMDA receptor antagonist, as a matter of convenience, it is preferred that these drugs be coadministered in a single dosage form. All modes of administrations are contemplated, e.g., orally, rectally, parenterally, topically, or by intravenous, intramuscular, intrasternal or subcutaneous injection or in a form suitable by inhalation. The formulations can, where appropriate, be conveniently presented in discrete dosage units and can be prepared by any of the methods well known in the art of pharmacy.

An analgesic composition containing the narcotic agonist-antagonist analgesic and nontoxic NMDA receptor antagonist will ordinarily be formulated with one or more pharmaceutically acceptable ingredients in accordance with known and established practice. Thus, the analgesic composition can be formulated as a liquid, powder, elixir, injectable solution, etc. Formulations for oral use can be provided as tablets or hard capsules wherein the pharmacologically active ingredients are mixed with an inert solid diluent such as calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredients are mixed with water or miscible solvents such as propylene glycol; PEG's and ethanol, or an oleaginous medium, e.g., peanut oil, liquid paraffin or olive oil.

For topical administration in the mouth, the analgesic compositions can take the form of buccal or sublingual tablets, drops or lozenges formulated in conventional manner.

For topical administration to the epidermis the compounds of the invention can be formulated as creams, gels, ointments or lotions or as transdermal patches. Such compositions can, for example, be formulated with an aqueous or oily base with the addition of suitable thickening, gelling, emulsifying, stabilizing, dispersing, suspending, and/or coloring agents.

The compounds of the invention can also be formulated as depot preparations. Such long acting formulations can be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example as a sparingly soluble salt.

The compounds of the invention can be formulated for parenteral administration by injection, conveniently intravenous, intramuscular or subcutaneous injection, for example by bolus injection or continuous intravenous

infusion. Formulations for injection can be presented in unit dosage from e.g. in ampoules or in multi-dose containers, with an added preservative. The compositions can take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

The compounds of the invention can also be formulated in rectal compositions such as suppositories or retention enemas, e.g. containing conventional suppository bases such as cocoa butter or other glyceride.

For intranasal administration, the compounds of the invention can be used, for example, as a liquid spray, as a powder or in the form of drops.

For administration by inhalation, the compounds according to the invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, tetrafluoroethane, heptafluoropropane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, for example, gelatin, for use in an inhaler or insulator can be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

Aqueous suspensions can include pharmaceutically acceptable excipients such as suspending agents, e.g., sodium carboxymethyl cellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as naturally occurring phosphatide, e.g., lecithin, or condensation products of an alkylene oxide with fatty acids, e.g., polyoxyethylene stearate, or condensation

products of ethylene oxide with long chain aliphatic alcohols, e.g., heptadecaethylene-oxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol, e.g., polyoxyethylene sorbitol monoleate or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, e.g., polyoxyethylene sorbitan monoleate. The aqueous suspensions can also contain one or more preservatives, e.g., ethyl-or-n-propyl-p-hydroxy benzoate, one or more coloring agents, one or more flavoring agents and one or more sweetening agents, such as sucrose, saccharin or sodium or calcium cyclamate.

In addition to the narcotic agonist-antagonist analgesic and nontoxic NMDA receptor antagonist, the analgesic composition herein can contain at least one other pharmacologically active substance, e.g., a non-narcotic analgesic such as acetaminophen, aspirin, diclofenac, diflusal, etodolac, fenbufen, fenoprofen, flufenisal, flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, meclofenamic acid, mefenamic acid, nabumetone, naproxen, oxaprozin, phenylbutazone, piroxicam, sulindac, tolmetin, tramadol, zomepirac, and the like.

EXAMPLES 1-39

The following unit dosage forms are illustrative of the pain-alleviating drug combinations in accordance with the present invention:

<u>Example</u>	<u>Narcotic Agonist-Antagonist Analgesic (mg)</u>	<u>Nontoxic NMDA Receptor Blocker (mg)</u>	<u>Additional Active Component (mg)</u>
1	pentazocine (35)	dextromethorphan hydrobromide (30)	
2	pentazocine (30)	dextromethorphan hydrobromide (30)	acetaminophen (325)
3	pentazocine (35)	dextrorphan	

5				
10			hydrobromide (30)	
	4	pentazocine (35)	amantadine (30)	
	5	pentazocine (35)	memantine (30)	
15	6	pentazocine (35)	memantine (30)	ibuprofen (325)
	7	nalbuphine (30)	dextromethorphan hydrobromide (30)	
20	8	nalbuphine (10)	dextromethorphan hydrobromide (30)	acetaminophen(325)
	9	nalbuphine (25)	dextrorphan hydrobromide (30)	
25				

	<u>Example</u>	<u>Narcotic Agonist-Antagonist Analgesic (mg)</u>	<u>Nontoxic NMDA Receptor Blocker (mg)</u>	<u>Additional Active Component (mg)</u>
5	10	nalbuphine (30)	amantadine (30)	
	11	nalbuphine (30)	memantine (30)	
10	12	nalbuphine (30)	memantine (30)	ibuprofen (325)
	13	butorphanol (10)	dextromethorphan hydrobromide (30)	
15	14	butorphanol (5)	dextromethorphan hydrobromide (30)	ibuprofen (325)
	15	butorphanol (10)	dextrophan hydrobromide (30)	
20	16	butorphanol (10)	amantadine (30)	
	17	butorphanol (10)	memantine	
25	18	butorphanol (10)	memantine	acetaminophen (325)
	19	buprenorphine (3)	dextromethorphan hydrobromide (30)	
30	20	buprenorphine (1)	dextromethorphan hydrobromide (30)	acetaminophen (325)
	21	buprenorphine (2)	dextrophan hydrobromide (30)	
35	22	buprenorphine (1)	amantadine (30)	
	23	buprenorphine (1)	memantine (30)	
	24	buprenorphine (1)	memantine (30)	ibuprofen (325)

	<u>Example</u>	<u>Narcotic Agonist-Antagonist Analgesic (mg)</u>	<u>Nontoxic NMDA Receptor Blocker (mg)</u>	<u>Additional Active Component (mg)</u>
5	25	meptazinol (30)	dextromethorphan hydrobromide (30)	
	26	meptazinol (35)	dextromethorphan hydrobromide (30)	acetaminophen (325)
10	27	meptazinol (30)	dextrorphan hydrobromide (30)	
	28	dezocine (30)	dextromethorphan hydrobromide (30)	
15	29	dezocine (25)	dextromethorphan hydrobromide (30)	acetaminophen (325)
20	30	dezocine (30)	dextrorphan hydrobromide (30)	
	31	dezocine (30)	amantadine (30)	
25	32	dezocine (30)	memantine (30)	
	33	dezocine (25)	memantine (30)	ibuprofen (325)
	34	nalorphine (35)	dextromethorphan hydrobromide (30)	
30	35	nalorphine (25)	dextromethorphan hydrobromide (30)	ibuprofen (325)

	<u>Example</u>	<u>Narcotic Agonist-Antagonist Analgesic (mg)</u>	<u>Nontoxic NMDA Receptor Blocker (mg)</u>	<u>Additional Active Component (mg)</u>
5	36	nalorphine (30)	dextrorphan hydrobromide (30)	
	37	cyclazocine (35)	dextromethorphan hydrobromide (30)	
10	38	cyclazocine (25)	dextromethorphan hydrobromide (30)	acetaminophen (325)
15	39	cyclazocine (30)	dextrorphan hydrobromide (30)	

In each of these dosage units, the nontoxic NMDA receptor antagonist dextromethorphan hydrobromide significantly potentiates the pain-alleviating activity of the narcotic agonist-antagonist analgesic.

20

WHAT IS CLAIMED IS:

1. An analgesic composition comprising (a) an analgesia-effective amount of at least one narcotic agonist-antagonist analgesic and (b) a narcotic agonist-antagonist analgesic-potentiating amount of at least one nontoxic N-methyl-D-aspartate receptor antagonist.
5
2. The analgesic composition of Claim 1 wherein the narcotic agonist-antagonist analgesic is selected from the group consisting of pentazocine, nalbuphine, butorphanol, buprenorphine, meptazinol, dezocine, nalorphine, cyclazocine and pharmaceutically acceptable salts thereof.
3. The analgesic composition of Claim 1 wherein the nontoxic NMDA receptor antagonist is at least one member selected from the group consisting of dextromethorphan, dextrorphan, amantadine, memantine and pharmaceutically acceptable salts thereof.
10
4. The analgesic composition of Claim 1 wherein (a) and (b) each is present in the same or different sustained release carrier.
15
5. The analgesic composition of Claim 1 containing a therapeutically effective amount of at least one other pharmaceutically active substance (c).
6. The analgesic composition of Claim 5 wherein the pharmacologically active substance (c) is a non-narcotic analgesic.
20
7. The analgesic composition of Claim 6 wherein the non-narcotic analgesic is selected from the group consisting of acetaminophen, aspirin, diclofenac, diflusal, etodolac, fenbufen, fenoprofen, flufenisal, flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, meclofenamic acid, mefenamic acid, nabumetone, naproxen, oxaprozin, phenylbutazone, piroxicam, sulindac, tolmetin, tramadol and zomepirac.
25
8. The analgesic composition of Claim 7 wherein the nontoxic NMDA receptor antagonist is selected from the group consisting of

dextromethorphan, dextrorphan, amantadine, memantine and pharmaceutically acceptable salts thereof.

9. The analgesic composition of Claim 1 containing at least 15 mg (b).
- 5 10. The analgesic composition of Claim 1 containing at least 20 mg (b).
- 10 11. A method of alleviating pain which comprises administering to a mammal exhibiting pain (a) an analgesia-effective amount of at least one narcotic agonist-antagonist analgesic and (b) a narcotic agonist-antagonist analgesic-potentiating amount of at least one nontoxic N-methyl-D-aspartate receptor antagonist.
- 15 12. The method of Claim 11 wherein the narcotic agonist-antagonist analgesic is selected from the group consisting of pentazocine, nalbuphine, butorphanol, buprenorphine, meptazinol, dezocine, nalorphine, cyclazocine and pharmaceutically acceptable salts thereof.
- 20 13. The method of Claim 11 wherein the nontoxic NMDA receptor antagonist is at least one member selected from the group consisting of dextromethorphan, dextrorphan, amantadine, memantine and pharmaceutically acceptable salts thereof.
- 25 14. The method of Claim 11 wherein (a) and (b) are coadministered.
15. The method of Claim 11 wherein (a) and (b) are coadministered as a sustained release dosage form.
16. The method of Claim 11 containing a therapeutically effective amount of at least one other pharmacologically active substance (c).
17. The method of Claim 16 wherein the pharmacologically active substance (c) is a non-narcotic analgesic.

18. The method of Claim 17 wherein the non-narcotic analgesic is selected from the group consisting of acetaminophen, aspirin, diclofenac, diflusal, etodolac, fenbufen, flufenisal, flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, meclofenamic acid, mefenamic acid, nabumetone, naproxen, oxaprozin, 5 phenylbutazone, piroxicam, sulindac, tolmetin, tramadol and zomepirac.

19. The method of Claim 18 wherein the nontoxic NMDA receptor antagonist is selected from the group consisting of dextromethorphan, dextrorphan, amantadine, memantine and pharmaceutically acceptable salts thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/05336

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K45/06 A61K31/485

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,Y	WO 97 04780 A (UNIV VIRGINIA COMMONWEALTH) 13 February 1997 * see in particular page 10 (examples 12,13); claims 1,15,20 *	1-19
Y	WIESENFELD-HALLIN Z: "Combined opioid-NMDA antagonist therapies. What advantages do they offer for the control of pain syndromes?." DRUGS, (1998 JAN) 55 (1) 1-4. REF: 37 JOURNAL CODE: EC2. ISSN: 0012-6667., XP002107488 New Zealand *see in particular p. 2, right column - page 3 *	1-19



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"S" document member of the same patent family

Date of the actual completion of the international search

28 June 1999

Date of mailing of the international search report

15/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Isert, B

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/05336

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9704780 A	13-02-1997	US 5840731 A	24-11-1998
		AU 6686596 A	26-02-1997
		CA 2228249 A	13-02-1997
		EP 0845989 A	10-06-1998
		US 5869498 A	09-02-1999
		US 5863922 A	26-01-1999
<hr/>			